



Concussion In Sports

WHAT YOU NEED TO KNOW



A Parent's Guide to Concussion in Sports

What is a concussion?

- A concussion is a brain injury which results in a temporary disruption of normal brain function. A concussion occurs when the brain is violently rocked back and forth or twisted inside the skull as a result of a blow to the head or body. An athlete does not have to lose consciousness ("knocked-out") to suffer a concussion.

Concussion Facts

- It is estimated that over 140,000 high school athletes across the United States suffer a concussion each year. (Data from NFHS Injury Surveillance System)
- Concussions occur most frequently in football, but girl's lacrosse, girl's soccer, boy's lacrosse, wrestling and girl's basketball follow closely behind. All athletes are at risk.
- A concussion is a traumatic injury to the brain.
- Concussion symptoms may last from a few days to several months.
- Concussions can cause symptoms which interfere with school, work, and social life.
- An athlete should not return to sports while still having symptoms from a concussion as they are at risk for prolonging symptoms and further injury.
- A concussion may cause multiple symptoms. Many symptoms appear immediately after the injury, while others may develop over the next several days or weeks. The symptoms may be subtle and are often difficult to fully recognize.

What are the signs and symptoms of a concussion?

SIGNS OBSERVED BY PARENTS, FRIENDS, TEACHERS OR COACHES	SYMPTOMS REPORTED BY ATHLETE
Appears dazed or stunned	Headache
Is confused about what to do	Nausea
Forgets plays	Balance problems or dizziness
Is unsure of game, score, or opponent	Double or fuzzy vision
Moves clumsily	Feeling sluggish
Answers questions slowly	Feeling foggy or groggy
Loses consciousness	Concentration or memory problems
Shows behavior or personality changes	Confusion
Can't recall events prior to hit	
Can't recall events after hit	



Concussion In Sports

WHAT YOU NEED TO KNOW



What should I do if I think my child has had a concussion?

If an athlete is suspected of having a concussion, he or she must be immediately removed from play, be it a game or practice. Continuing to participate in physical activity after a concussion can lead to worsening concussion symptoms, increased risk for further injury, and even death. Parents and coaches are not expected to be able to “diagnose” a concussion, as that is the job of a medical professional. However, you must be aware of the signs and symptoms of a concussion and if you are suspicious, then your child must stop playing:

When in doubt, sit them out!

All athletes who sustain a concussion need to be evaluated by a health care professional who is familiar with sports concussions. You should call your child’s physician and explain what has happened and follow your physician’s instructions. If your child is vomiting, has a severe headache, is having difficulty staying awake or answering simple questions he or she should be taken to the emergency department immediately.

When can an athlete return to play following a concussion?

After suffering a concussion, **no athlete should return to play or practice on that same day.** Previously, athletes were allowed to return to play if their symptoms resolved within 15 minutes of the injury. Studies have shown us that the young brain does not recover quickly enough for an athlete to return to activity in such a short time.

Concerns over athletes returning to play too quickly have led state lawmakers in both Oregon and Washington to pass laws stating that **no player shall return to play following a concussion on that same day and the athlete must be cleared by an appropriate health-care professional before he or she are allowed to return to play in games or practices.** The laws also mandate that coaches receive education on recognizing the signs and symptoms of concussion.

Once an athlete no longer has symptoms of a concussion and is cleared to return to play by health care professional knowledgeable in the care of sports concussions he or she should proceed with activity in a step-wise fashion to allow the brain to re-adjust to exertion. On average the athlete will complete a new step each day. The return to play schedule should proceed as below following medical clearance:

Step 1: Light exercise, including walking or riding an exercise bike. No weight-lifting.

Step 2: Running in the gym or on the field. No helmet or other equipment.

Step 3: Non-contact training drills in full equipment. Weight-training can begin.

Step 4: Full contact practice or training.

Step 5: Game play.

If symptoms occur at any step, the athlete should cease activity and be re-evaluated by their health care provider.



Concussion In Sports

WHAT YOU NEED TO KNOW



How can a concussion affect schoolwork?

Following a concussion, many athletes will have difficulty in school. These problems may last from days to months and often involve difficulties with short and long-term memory, concentration, and organization.

In many cases it is best to lessen the athlete's class load early on after the injury. This may include staying home from school for a few days, followed by a lightened schedule for a few days, or perhaps a longer period of time, if needed. Decreasing the stress on the brain early on after a concussion may lessen symptoms and shorten the recovery time.

What can I do?

- Both you and your child should learn to recognize the "Signs and Symptoms" of concussion as listed above.
- Teach your child to tell the coaching staff if he or she experiences such symptoms.
- Emphasize to administrators, coaches, teachers, and other parents your concerns and expectations about concussion and safe play.
- Teach your child to tell the coaching staff if he or she suspects that a teammate has a concussion.
- Monitor sports equipment for safety, fit, and maintenance.
- Ask teachers to monitor any decrease in grades or changes in behavior that could indicate concussion.
- Report concussions that occurred during the school year to appropriate school staff. This will help in monitoring injured athletes as they move to the next season's sports.

Other Frequently Asked Questions

Why is it so important that an athlete not return to play until they have completely recovered from a concussion?

Athletes who are not fully recovered from an initial concussion are significantly vulnerable for recurrent, cumulative, and even catastrophic consequences of a second concussive injury. Such difficulties are prevented if the athlete is allowed time to recover from the concussion and return to play decisions are carefully made. No athlete should return-to-sport or other at-risk participation when symptoms of concussion are present and recovery is ongoing.

Is a "CAT scan" or MRI needed to diagnose a concussion?

Diagnostic testing, which includes CT ("CAT") and MRI scans, are rarely needed following a concussion. While these are helpful in identifying life-threatening brain injuries (e.g. skull fracture, bleeding, swelling), they are not normally utilized, even by athletes who have sustained severe concussions. A concussion is diagnosed based upon the athlete's story of the injury and the health care provider's physical examination.



Concussion In Sports

WHAT YOU NEED TO KNOW



What is the best treatment to help my child recover more quickly from a concussion?

The best treatment for a concussion is rest. There are no medications that can speed the recovery from a concussion. Exposure to loud noises, bright lights, computers, video games, television and phones (including text messaging) all may worsen the symptoms of a concussion. You should allow your child to rest as much as possible in the days following a concussion. As the symptoms lessen, you can allow increased use of computers, phone, video games, etc., but the access must be lessened if symptoms worsen.

How long do the symptoms of a concussion usually last?

The symptoms of a concussion will usually go away within one week of the initial injury. You should anticipate that your child will likely be out of sports for about two weeks following a concussion. However, in some cases symptoms may last for several weeks, or even months. Symptoms such as headache, memory problems, poor concentration, and mood changes can interfere with school, work, and social interactions. The potential for such long-term symptoms indicates the need for careful management of all concussions.

How many concussions can an athlete have before he or she should stop playing sports?

There is no “magic number” of concussions that determine when an athlete should give up playing contact or collision sports. The circumstances surrounding each individual injury, such as how the injury happened and length of symptoms following the concussion, are very important and must be considered when assessing an athlete’s risk for further and potentially more serious concussions. The decision to “retire” from sports is a decision best reached following a complete evaluation by your child’s primary care provider and consultation with a physician or neuropsychologist who specializes in treating sports concussion.

I’ve read recently that concussions may cause long-term brain damage in professional football players. Is this a risk for high school athletes who have had a concussion?

The issue of “chronic encephalopathy” in several former NFL players has received a great deal of media attention lately. Very little is known about what may be causing dramatic abnormalities in the brains of these unfortunate retired football players. At this time we have very little knowledge of the long-term effects of concussions which happen during high school athletics.

In the cases of the retired NFL players, it appears that most had long careers in the NFL after playing in high school and college. In most cases, they played football for over 20 years and suffered multiple concussions in addition to hundreds of other blows to their heads. Alcohol and steroid use may also be contributing factors in some cases. Obviously, the average high school athlete does not come close to suffering the total number or shear force of head trauma seen by professional football players. However, the fact that we know very little about the long-term effects of concussions in young athletes is further reason to very carefully manage each concussion.



Concussion In Sports

WHAT YOU NEED TO KNOW



Some of this information has been adapted from the CDC's "Heads Up: Concussion in High School Sports" materials by the NFHS's Sports Medicine Advisory Committee. Please go to www.cdc.gov/ncipc/tbi/Coaches_Tool_Kit.htm for more information.

If you have any further questions regarding concussions in high school athletes or want to know how to find a concussion specialist in your area please contact Michael C. Koester, MD, ATC and Chair of the NFHS Sports Medicine Advisory Committee at michael.koester@slocumcenter.com.

April 2010

HEADS+UP

CONCUSSION IN HIGH SCHOOL SPORTS

A FACT SHEET FOR **ATHLETES**

What is a concussion?

A concussion is a brain injury that:

- Is caused by a bump, blow, or jolt to the head or body.
- Can change the way your brain normally works.
- Can occur during practices or games in any sport or recreational activity.
- Can happen even if you haven't been knocked out.
- Can be serious even if you've just been "dinged" or "had your bell rung."

All concussions are serious. A concussion can affect your ability to do schoolwork and other activities (such as playing video games, working on a computer, studying, driving, or exercising). Most people with a concussion get better, but it is important to give your brain time to heal.

What are the symptoms of a concussion?

You can't see a concussion, but you might notice **one or more** of the symptoms listed below or that you "don't feel right" soon after, a few days after, or even weeks after the injury.

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Bothered by light or noise
- Feeling sluggish, hazy, foggy, or groggy
- Difficulty paying attention
- Memory problems
- Confusion

What should I do if I think I have a concussion?

- **Tell your coaches and your parents.** Never ignore a bump or blow to the head even if you feel fine. Also, tell your coach right away if you think you have a concussion or if one of your teammates might have a concussion.
- **Get a medical check-up.** A doctor or other health care professional can tell if you have a concussion and when it is OK to return to play.
- **Give yourself time to get better.** If you have a concussion, your brain needs time to heal. While your brain is still healing, you are much more likely to have another concussion. Repeat concussions can increase the time it takes for you to recover and may cause more damage to your brain. It is important to rest and not return to play until you get the OK from your health care professional that you are symptom-free.

How can I prevent a concussion?

Every sport is different, but there are steps you can take to protect yourself.

- Use the proper sports equipment, including personal protective equipment. In order for equipment to protect you, it must be:
 - The right equipment for the game, position, or activity
 - Worn correctly and the correct size and fit
 - Used every time you play or practice
- Follow your coach's rules for safety and the rules of the sport.
- Practice good sportsmanship at all times.

If you think you have a concussion:
Don't hide it. Report it. Take time to recover.

It's better to miss one game than the whole season.

For more information and to order additional materials *free-of-charge*, visit: www.cdc.gov/Concussion.

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION



ATENCIÓN*

CONMOCIONES CEREBRALES EN LOS DEPORTES DE LA ESCUELA SECUNDARIA

HOJA INFORMATIVA PARA **LOS ATLETAS**

¿Qué es una conmoción cerebral?

Una conmoción cerebral es una lesión del cerebro que:

- Es causada por un golpe o una sacudida en la cabeza o el cuerpo.
- Puede alterar el funcionamiento normal del cerebro.
- Puede ocurrir durante las prácticas o la competición de cualquier deporte o durante las actividades recreativas.
- Puede ocurrir aun cuando no se haya perdido el conocimiento.
- Puede ser grave aunque se trate de un golpe leve o que provoque una sensación de zumbido en la cabeza.

Todas las conmociones cerebrales son graves. Las conmociones cerebrales pueden afectar tus actividades escolares u otras actividades (como jugar video juegos, trabajar en la computadora, estudiar, conducir o hacer ejercicio). La mayoría de las personas que sufren una conmoción cerebral se mejoran, pero es importante tomarse el tiempo necesario para que el cerebro se recupere.

¿Cuáles son los síntomas de una conmoción cerebral?

Aunque la conmoción cerebral no se pueda observar, puede que notes uno o más de los siguientes síntomas o que “no te sientas del todo bien” justo después de la lesión, a los días o las semanas siguientes.

- Dolor de cabeza o “presión” en la cabeza
- Náuseas o vómitos
- Problemas de equilibrio o mareo
- Visión borrosa o doble
- Molestia causada por la luz o el ruido
- Debilidad, confusión, aturdimiento o estado grogui
- Dificultad para prestar atención
- Problemas de memoria
- Confusión

¿Qué debo hacer si creo que he sufrido una conmoción cerebral?

- Avísale a tus entrenadores y a tus padres. Nunca ignores un golpe o una sacudida en la cabeza, aun cuando te sientas bien. También, avísale a tu entrenador enseguida si crees que has sufrido una conmoción cerebral o le puede haber pasado a uno de tus compañeros.
- Ve al médico para que te examine. Un médico u otro profesional de la salud podrá decirte si sufriste una conmoción cerebral y cuándo estarás listo para volver a jugar.
- Tómate el tiempo suficiente para curarte. Si sufriste una conmoción cerebral, tu cerebro necesitará tiempo para sanarse. Cuando tu cerebro se está curando, existe una mayor probabilidad de que sufras una segunda conmoción. Las conmociones cerebrales repetidas pueden aumentar el tiempo de recuperación y dañar más el cerebro. Es importante descansar y no volver a jugar hasta que tu profesional de la salud te indique que ya no tienes más síntomas y que puedes reanudar tu actividad deportiva.

¿Cómo puedo prevenir una conmoción cerebral?

Depende del deporte que practicas, pero puedes tomar una serie de medidas para protegerte.

- Usa el equipo de deporte adecuado, incluido el equipo de protección personal. Para que este equipo te proteja, debe:
 - Ser adecuado para el deporte que practicas, tu posición en el juego y tipo de actividad.
 - Usarse correctamente y ajustarse bien a tu cuerpo.
 - Colocarse cada vez que juegues o practiques.
- Sigue las reglas de seguridad del entrenador y las reglas del deporte que practicas.
- Mantén el espíritu deportivo en todo momento.

Si crees que sufriste una conmoción cerebral:
No trates de ocultarlo. Notifícaselo a alguien.
Tómate tiempo para recuperarte.

Es preferible perderse un juego que toda la temporada.

Para obtener más información y solicitar más materiales *de forma gratuita*, visite: www.cdc.gov/Concussion.

DEPARTAMENTO DE SALUD Y SERVICIOS HUMANOS DE LOS EE. UU.
CENTROS PARA EL CONTROL Y LA PREVENCIÓN DE ENFERMEDADES

